

COORBITS FOR PROJECTIVE REPRESENTATIONS WITH AN APPLICATION TO BERGMAN SPACES

JENS GERLACH CHRISTENSEN, AMER H. DARWEESH, AND GESTUR ÓLAFSSON

ABSTRACT. Recently representation theory has been used to provide atomic decompositions for a large collection of classical Banach spaces. In this paper we extend the techniques to also include projective representations. As our main application we obtain atomic decompositions of Bergman spaces on the unit ball through the holomorphic discrete series for the group of isometries of the ball.

1. INTRODUCTION

With the rise of continuous wavelet theory it was discovered that representation theory could be used to obtain atomic decompositions for some classical Banach spaces. This area of harmonic analysis is called coorbit theory and it was initiated by Feichtinger and Gröchenig [19, 20, 18, 22]. Several interesting generalizations were later presented in [9, 28, 21, 29, 13, 14, 15, 12]. All these examples use irreducible integrable representations in order to construct atomic decompositions. This allows one to choose atoms in an appropriate minimal Banach space. As has been remarked recently, the restriction of irreducibility and integrability as well as the criteria for selecting atoms turns out to be too restrictive. Therefore, the first and last author suggested the use of Fréchet spaces [7, 5] in coorbit theory, and recently the idea has been used in several cases [6, 8, 4, 11].

There are several situations where it is more natural to consider projective representations rather than representations. Typical examples are the Bergman spaces on bounded symmetric domains. This paper is concerned with extending the theory in [7] and [5] to projective representations. The idea of using projective representations in coorbit theory has earlier been explored in [9] under the assumption of irreducibility and integrability. Those two restrictions were removed in the thesis of the second author [16]. Upon closer inspection the theories presented in [9] and [16] both generally only apply to simply connected groups. The reason is that the multiplier (or cocycle) for the projective representation is assumed continuous, and this property can only be ensured for simply connected groups. In particular [9] and [16] cannot be used to describe Bergman spaces on the unit ball in $\mathbb{C}^n$ as coorbits for the group $SU(n, 1)$ which is not simply connected. However, for general locally compact groups it can be proven that the multiplier can be chosen continuous in a neighborhood of the identity, and for Lie groups the multiplier can even be chosen smooth in a neighborhood of the identity. In this paper we show that

The research was partially supported by NSF grant DMS 1321794. The first and last named authors would also like to thank AMS for it's support during the MRC program *Lie Group Representations, Discretization, and Gelfand Pairs* June 5–June 11, 2016 .

these observations are sufficient for obtaining a working coorbit theory for projective representations and for finding atomic decompositions of Bergman spaces on the unit ball. We thereby finish a study initiated in [4] which vastly increases the class of atoms used to decompose Bergman spaces when compared to the classical results in [10, 25]. In [4] we were restricted to using discrete series representations of $SU(n, 1)$ with a rational parameter, and therefore we were not able to cover the entire scale of Bergman spaces. The use of projective representations allows us to also work with representations with an irrational parameter and this is used to close the gap in [4].

2. PROJECTIVE REPRESENTATIONS

In this section we review known results on measurable and locally continuous projective representations, and, following Mackey, we will construct a representation of an extension of a locally compact group from a given projective representation. We use [31] as a standard reference even if the results are mostly due to Mackey and Bargmann.

We assume that G is a locally compact group equipped with a fixed left invariant Haar measure which we denote by dx . We denote by $\mathbb{T} := \{t \in \mathbb{C} \mid |t| = 1\}$ the one dimensional torus with normalized Haar measure dt .

Definition 2.1. Let $\mathcal{S}$ be a locally convex Hausdorff topological vector space over $\mathbb{C}$, and denote by $\mathcal{S}^*$ its conjugate dual. A projective representation of G is a mapping $\rho : G \rightarrow GL(\mathcal{S})$, the space of continuous linear maps $\mathcal{S} \rightarrow \mathcal{S}$ with a continuous inverse, that satisfies the following three conditions:

- (1) $\rho(1) = \text{id}$.
- (2) There is a Borel function (called a *multiplier* or a *cocycle*) $\sigma : G \times G \rightarrow \mathbb{T}$, which satisfies the condition

$$\rho(xy) = \sigma(x, y)\rho(x)\rho(y).$$

- (3) For every $v \in \mathcal{S}$ and every $\lambda \in \mathcal{S}^*$ the mapping

$$x \mapsto \langle \lambda, \rho(x)v \rangle := \lambda(\rho(x)v)$$

is a Borel function.

A projective representation with multiplier σ is said to be a σ -representation.

Notice, that for a σ -representation ρ and $x \in G$ we have

$$\rho(x^{-1}) = \overline{\sigma(x, x^{-1})}\rho(x)^{-1} = \overline{\sigma(x^{-1}, x)}\rho(x)^{-1}. \quad (2.1)$$

Let $x, y, z \in G$. The following are straightforward consequences about the cocycle σ :

- (1) $\sigma(x, 1) = \sigma(1, x) = 1$,
- (2) $\sigma(x, y)^{-1} = \sigma(x, y)$,
- (3) $\sigma(x, x^{-1}) = \sigma(x^{-1}, x)$,
- (4) $\sigma(xy, z)\sigma(x, y) = \sigma(x, yz)\sigma(y, z)$.

Following Bargmann, we say that two cocycles σ and τ are *similar* if there exists a Borel function $a : G \rightarrow \mathbb{T}$ such that

$$\tau(x, y) = \frac{a(xy)}{a(x)a(y)}\sigma(x, y). \quad (2.2)$$

We note that if τ is similar to σ via the Borel function a and ρ is a σ -representation, then $\eta(x) := a(x)\rho(x)$ is a τ -representations.

Theorem 2.2. *Every multiplier is similar to a multiplier which is continuous on some open neighborhood of $(e, e) \in G \times G$. If G is a Lie group, then every multiplier is similar to a multiplier that is analytic in an open neighborhood of (e, e) . If G is a connected and simply connected Lie group, then every multiplier is similar to a multiplier which is analytic on the entire group $G \times G$.*

Proof. The first statement is [31, Corollary 7.6]. The second statement is [31, Corollary 7.21] and the last segment is [31, Corollary 7.30]. $\square$

For projective representations we define irreducibility, cyclicity, admissible vectors, unitarity and square integrability in the same way as for representations. We summarize these notions in the following definition.

Definition 2.3. Let $(\rho, \mathcal{S})$ be a projective representation, then

- (1) A subspace W of $\mathcal{S}$ is ρ -invariant if $\rho(x)W \subseteq W$ for all $x \in G$.
- (2) $(\rho, \mathcal{S})$ is irreducible if the only closed ρ -invariant subspaces are $\{0\}$ and $\mathcal{S}$ itself.
- (3) A vector $u \in \mathcal{S}$ is a ρ -cyclic if $\text{span}\{\rho(x)u \mid x \in G\}$ is dense in $\mathcal{S}$. If such a cyclic vector exists, we say that $(\rho, \mathcal{S})$ is a cyclic projective representation.
- (4) $(\rho, \mathcal{H})$ is unitary if $\mathcal{H}$ is a Hilbert space, and $\rho(x)$ is a unitary operator for every $x \in G$.
- (5) If $\mathcal{H}$ is a Hilbert space and $(\rho, \mathcal{H})$ is irreducible and unitary it is called square-integrable if there is a nonzero vector $u \in \mathcal{H}$ such that

$$\int_G |\langle u, \rho(x)u \rangle|^2 dx < \infty.$$

In this case u is called a ρ -admissible vector.

In the following lemma we define the dual projective representation on the conjugate dual of a Fréchet space.

Lemma 2.4. *Let $(\rho, \mathcal{S})$ be a σ -representation of G on a Fréchet space $\mathcal{S}$, and let $\mathcal{S}^*$ be the conjugate dual of $\mathcal{S}$ equipped with the weak*-topology. The mapping ρ^* , which is given by*

$$\langle \rho^*(x)\lambda, v \rangle := \langle \lambda, \rho(x)^{-1}v \rangle$$

for all $\lambda \in \mathcal{S}^*$ and all $v \in \mathcal{S}$, defines a σ -representation of G on the space $\mathcal{S}^*$. Finally $x \mapsto \langle \rho^*(x)\lambda, u \rangle$ is continuous around e if σ and ρ are locally continuous.

Proof. As $\mathcal{S}^*$ is equipped with the weak* topology, its conjugate dual is $\mathcal{S}$. We denote this dual pairing by $\langle\langle \cdot, \cdot \rangle\rangle$. For $v \in (\mathcal{S}^*)^* = \mathcal{S}$ and $\lambda \in \mathcal{S}^*$ we have $\langle\langle v, \lambda \rangle\rangle = \overline{\langle \lambda, v \rangle}$. Therefore the following calculation shows that ρ^* has cocycle σ

$$\begin{aligned} \langle \rho^*(xy)\lambda, v \rangle &= \langle \lambda, \rho(xy)^{-1}v \rangle \\ &= \langle \lambda, (\sigma(x, y)\rho(x)\rho(y))^{-1}v \rangle \\ &= \langle \lambda, \overline{\sigma(x, y)}\rho(y)^{-1}\rho(x)^{-1}v \rangle \\ &= \langle \sigma(x, y)\lambda, \rho(y)^{-1}\rho(x)^{-1}v \rangle \\ &= \langle \sigma(x, y)\rho^*(x)\rho^*(y)\lambda, v \rangle \end{aligned}$$

Hence, $\rho^*(xy) = \sigma(x, y)\rho^*(x)\rho^*(y)$. Furthermore, the equation (2.1) implies that

$$\langle\langle v, \rho^*(x)\lambda \rangle\rangle = \overline{\langle \rho^*(x)\lambda, v \rangle} = \overline{\langle \lambda, \rho(x)^{-1}u \rangle} = \sigma(x^{-1}, x)\overline{\langle \lambda, \rho(x^{-1})u \rangle}.$$

Since $x \mapsto x^{-1}$ is continuous and σ and ρ are Borel it follows that $x \mapsto \langle\langle v, \rho^*(x)\lambda \rangle\rangle$ is Borel. Moreover, the mapping is locally continuous if σ and ρ are locally continuous. $\square$

For any projective representation ρ of G we can construct an actual representation of a new group related to G which is called the Mackey obstruction group of G (see p. 269f in [26]). We refer to [31, Chap. VII] for a detailed discussion.

We first gather some facts about the Mackey group. Let σ be a cocycle on $G \times G$. As a set, the Mackey group that corresponds to G is the group $G_\sigma := G \times \mathbb{T}$, with multiplication given by

$$(x, t)(y, s) = (xy, \overline{\sigma(x, y)}ts).$$

The inverse of $(x, t) \in G_\sigma$ is given by

$$(x, t)^{-1} = (x^{-1}, \sigma(x, x^{-1})\bar{t}) = (x^{-1}, \sigma(x^{-1}, x)\bar{t}).$$

Note that this is a *central* extension of $\mathbb{T}$ by G as all the elements (e, s) , $s \in \mathbb{T}$, are central in G_σ . The product of the Borel algebras of G and $\mathbb{T}$ defines a σ -algebra on G_σ and the product measure $dxd t$ is left invariant.

Theorem 2.5. *Let the notation be as above. Then the following holds:*

- (1) *There exists a unique topology on G_σ , called the Weyl topology, that generates the product σ -algebra and at the same time makes G_σ into a locally compact Hausdorff topological group.*
- (2) *If the multiplier σ is continuous around (e, e) , then there exists a neighborhood U of e in G such that the Weyl topology on $U \times \mathbb{T}$ corresponds to the product topology on $U \times \mathbb{T}$.*
- (3) *Two extensions G_σ and G_τ are isomorphic if and only if the multipliers σ and τ are similar. If the similarity is given by the function a as in (2.2) then the isomorphism $G_\sigma \rightarrow G_\tau$ is given by $(x, t) \mapsto (x, \overline{a(x)}t)$.*
- (4) *If G is a connected Lie group, then there exists a unique Lie group structure on G_σ compatible with the Weyl topology. The maps $t \mapsto (e, t)$ and $(x, t) \mapsto x$ are analytic.*
- (5) *There exists a smooth, respectively analytic, multiplier τ similar to σ if and only if there exists a smooth, respectively analytic, map $\kappa : G \rightarrow G_\sigma$ such that $p_1(\kappa(x)) = x$ where $p_1(x, t) = x$.*

Proof. (1) and (3) are [31, Theorem 7.8]. (2) is [31, Corollary 7.10]. (4) is [31, Theorem 7.21]. Finally, (5) is [31, Corollary 7.23]. $\square$

Next we discuss the construction of a representation of G_σ from a σ -representation ρ . Define

$$\rho_\sigma(x, t) := t\rho(x).$$

Then a simple calculation shows that ρ_σ is a homomorphism. If τ is similar to σ , $\eta(x) = \overline{a(x)}\rho(x)$ is the corresponding canonical τ -representation and $\varphi(x, t) = \overline{a(x)}\rho(x)$ is the natural isomorphism $G_\sigma \simeq G_\tau$, then

$$\eta_\tau(\varphi(x, t)) = \overline{a(x)}t\eta(x) = \overline{a(x)}a(x)t\rho(x) = \rho_\sigma(x, t)$$

as $a(x) \in \mathbb{T}$. Hence $\eta_\tau \circ \varphi = \rho_\sigma$.

The following is well known for representations on a separable Hilbert space, see [31, Theorem 7.16].

Theorem 2.6. *Assume that the multiplier σ is continuous in a neighborhood of (e, e) , and that G is connected. Then*

$$G_\sigma \rightarrow \mathbb{C}, \quad (x, t) \mapsto \langle \lambda, \rho_\sigma(x, t)u \rangle = \bar{t} \langle \lambda, \rho(x)u \rangle \quad (2.3)$$

is a Borel function for all $\lambda \in \mathcal{S}^*$ and $u \in \mathcal{S}$. Furthermore the following holds:

- (1) *The map in (2.3) is continuous if and only if $G \rightarrow \mathbb{C}, x \mapsto \langle \lambda, \rho(x)u \rangle$ is continuous in an open neighborhood around e .*
- (2) *Assume that G is a Lie group. The map in (2.3) is smooth if and only if $G \rightarrow \mathbb{C}, x \mapsto \langle \lambda, \rho(x)u \rangle$ is smooth in an open neighborhood around e .*

Proof. As σ is continuous around (e, e) there exists, according to Theorem 2.5, part (2), a e -neighborhood U in G such that the Weyl topology on $U \times \mathbb{T}$ agrees with the product topology. Thus the map in (2.3) is Borel on $U \times \mathbb{T}$. Let $z = (y, s) \in G_\sigma$. Then $U \times \mathbb{T}(y, 1)$ is a neighborhood around z and for $(x, t) \in U \times \mathbb{T}$ we have

$$(x, t) \mapsto \lambda(\rho_\sigma((x, t)(y, 1))u) = \lambda(\rho_\sigma((x, t))\rho_\sigma((y, 1))u)$$

which is Borel in (x, t) . Hence the map in (2.3) is Borel. The same argument proves that (2.3) is continuous if $x \mapsto \langle \lambda, \rho(x)u \rangle$ is locally continuous. Let us check the opposite direction and assume that the mapping (2.3) is continuous. Restrict the mapping to a neighbourhood $U \times \mathbb{T}$ in the Weyl topology for which U is open in G . The mapping from U to $U \times \mathbb{T}$ given by $x \rightarrow (x, 1)$ is then continuous, and therefore $x \mapsto \langle \lambda, \rho_\sigma(x, 1)u \rangle = \langle \lambda, \rho(x)u \rangle$ is continuous on the neighbourhood U .

Smoothness is verified in the same manner. $\square$

Assumption 2.7. From now on we will assume that multipliers are continuous in a neighborhood of (e, e) . Moreover, we will assume that for every $\lambda \in \mathcal{S}^*$ and every $u \in \mathcal{S}$, there is a neighbourhood U of e on which the mapping $x \mapsto \langle \lambda, \rho(x)u \rangle$ is continuous. Finally the group G is assumed to be connected.

The following examples show that there are plenty of examples for which this assumption is satisfied.

Example 2.8. If $\mathcal{H}$ is a Hilbert space and ρ is a unitary projective representation, then Corollary 7.10 in [31] ensures that there is a neighbourhood U on which $x \mapsto \langle \lambda, \rho(x)u \rangle$ is continuous.

We also note that the representation ρ_σ is unitary if and only if the σ -projective representation ρ is unitary. Finally, the σ -representation ρ is square-integrable if and only if ρ_σ -is square integrable. This last statement follows from the fact that $(u, \rho_\sigma(x, t)u) = \bar{t}(u, \rho(x)u)$ and that $\mathbb{T}$ is compact with measure 1. Thus $\int_{G_\sigma} |(u, \rho_\sigma(x, t)u)|^2 dx dt = \int_G |(u, \rho(x)u)|^2 dx$

We now provide a version of the Duflo-Moore theorem for square integrable projective representations. This result can be found in [1].

Theorem 2.9. *Let $(\rho, \mathcal{H})$ be a square-integrable projective representation of G .*

- (1) *There exists a positive self adjoint operator A_ρ which is defined on a dense subset D of $\mathcal{H}$, such that $u \in \mathcal{H}$ is ρ -admissible if and only if $u \in D$. Moreover, the orthogonality relation*

$$\int_G (v_1, \rho(x)u_1) (\rho(x)u_2, v_2) dx = (A_\rho u_2, A_\rho u_1) (v_1, v_2)$$

holds for all $u_1, u_2 \in D$ and $v_1, v_2 \in \mathcal{H}$.

- (2) In addition, if G is a unimodular, then $D = \mathcal{H}$ and $A_\rho = c_\rho \text{Id}_{\mathcal{H}}$. Thus, all vectors of $\mathcal{H}$ are ρ -admissible and

$$\int_G (v_1, \rho(x)u_1) (\rho(x)u_2, v_2) dx = c_\rho^2(u_2, u_1) (v_1, v_2)$$

for all $u_1, u_2, v_1, v_2 \in \mathcal{H}$. The constant $1/c_\rho^2$ is called the formal dimension of ρ .

Example 2.10. We will now show that for a specific Gelfand triple $(\mathcal{S}, \mathcal{H}, \mathcal{S}^*)$ from the original coorbit theory [18, 19, 20, 9] the weak continuity requirement from Assumption 2.7 is automatically satisfied.

Let $(\rho, \mathcal{H})$ be a unitary σ -representation of G , and let ρ_σ be the Mackey representation of G_σ . Then $(\rho_\sigma, \mathcal{H})$ is a (strongly) continuous representation (follows from Example 2.8 and Theorem 2.6). Moreover, assume that ρ is an irreducible square integrable projective representation and assume there is a non-zero u for which $(u, \rho(\cdot)u)$ is in $L_w^1(G)$ for some submultiplicative weight $w \geq 1$, see Remark 3.2 below for definition. Let $\mathcal{S}$ be the Banach space

$$\mathcal{S} = \mathcal{H}_w^1 = \{v \in \mathcal{H} \mid (v, \rho(\cdot)u) \in L_w^1(G)\}$$

equipped with the norm it inherits from $L_w^1(G)$. It is clear that for a cocycle σ for ρ the space $\mathcal{H}_w^1$ is isometrically isomorphic to the space

$$\{v \in \mathcal{H} \mid (v, \rho_\sigma(\cdot)u) \in L_w^1(G_\sigma)\}$$

when $w(x, t) = w(x)$.

From standard coorbit theory [19] the representation ρ_σ restricted to H_w^1 is strongly continuous (because left translation is continuous on $L_w^1(G_\sigma)$). Therefore it follows immediately that $(x, t) \mapsto \langle \lambda, \rho_\sigma(x, t)v \rangle$ is continuous on G_σ for $\lambda \in (H_w^1)^*$ and $v \in H_w^1$. Then from Theorem 2.6 it follows that $x \mapsto \langle \lambda, \rho(x)v \rangle$ is continuous on a neighbourhood of e .

Example 2.11. Now assume that G is a Lie group. We will now show that the smooth vectors for a projective representation satisfy the weak continuity from Assumption 2.7.

The smooth vectors for the projective representation ρ on the Hilbert space $\mathcal{H}$ is the collection of vectors u for which $G \ni x \mapsto \rho(x)u \in \mathcal{H}$ is C^∞ on a neighbourhood of e . By arguments similar to those in [27] this is the equivalent to the weak smoothness of the mapping $x \mapsto (v, \rho(x)u)$ for any $v \in \mathcal{H}$ on a neighbourhood of e . By Theorem 2.6 we then get that $(x, t) \mapsto (v, \rho_\sigma(x, t)u)$ is smooth, which tells us that u is a smooth vector for ρ_σ by [27]. This argument also works in the reverse direction, so we see that the smooth vectors for ρ are the same as the smooth vectors for ρ_σ , i.e. $\mathcal{H}_\rho^\infty = \mathcal{H}_{\rho_\sigma}^\infty$. Since $\rho_\sigma(x, t)u = t\rho(x)u$ the derivatives in t are just multiples of the identity. Therefore the usual Fréchet space topologies on $\mathcal{H}_\rho^\infty$ and $\mathcal{H}_{\rho_\sigma}^\infty$ are generated by the same differential operators, and therefore their topologies are equivalent. Therefore they also have the same dual spaces $\mathcal{H}_\rho^{-\infty} = \mathcal{H}_{\rho_\sigma}^{-\infty}$.

It is classical, see [33], that ρ_σ restricted to $\mathcal{H}_{\rho_\sigma}$ is a continuous representation. Therefore $(x, t) \mapsto \langle \lambda, \rho_\sigma(x, t)u \rangle$ is continuous on G_σ for $\lambda \in \mathcal{H}_{\rho_\sigma}^{-\infty}$ and $u \in \mathcal{H}_{\rho_\sigma}^\infty$. Therefore Theorem 2.6 ensures that $x \mapsto \langle \lambda, \rho(x)u \rangle$ is continuous on a neighbourhood of e .

Let $u \in \mathcal{S}$ be ρ -cyclic vector. We define the wavelet transform $W_u^\rho : \mathcal{S}^* \rightarrow C(G)$ by

$$W_u^\rho(\lambda)(x) := \langle \lambda, \rho(x)u \rangle.$$

In the following lemma we state the relation between a projective representation and the corresponding representation of the Mackey group. The proof is straightforward from the definition.

Lemma 2.12. *Let $(\rho, \mathcal{S})$ be a projective representation of G and let $(\rho_\sigma, \mathcal{S})$ be the corresponding representation of G_σ . Then the following are true:*

- (1) *The vector $u \in \mathcal{S}$ is ρ -cyclic if and only if u is ρ_σ -cyclic.*
- (2) *The wavelet transforms are related by $W_u^{\rho_\sigma}(\lambda)(x, t) = t W_u^\rho(\lambda)(x)$.*

3. BANACH FUNCTION SPACES AND SEQUENCE SPACE

In this section we define twisted translation and twisted convolution, and we summarize the assumptions we will place on a Banach space of functions (or BF-space for short) throughout this paper. We will also introduce a collection of sequence spaces which will be used in Sections 5 and 6 in order to formulate our results on atomic decompositions.

Definition 3.1. Let B be a Banach space over $\mathbb{C}$ of functions on G , and let σ be a cocycle on G . For a function $f \in B$, we define twisted left translation by

$$\ell_y^\sigma f(x) := \overline{\sigma(y, y^{-1}x)} f(y^{-1}x),$$

and we define twisted right translation by

$$r_y^\sigma f(x) := \sigma(x, y) f(xy).$$

If the cocycle is 1 we set $\ell_y = \ell_y^\sigma$ and $r_y = r_y^\sigma$, and we drop the use of the word twisted.

It is important to notice the following relations between the translation operators and the wavelet transform

$$\begin{aligned} \ell_y^\sigma W_u^\rho(\lambda)(x) &= W_u^\rho(\rho^*(y)\lambda)(x) \\ r_y^\sigma W_u^\rho(\lambda)(x) &= W_{\rho(y)u}^\rho(\lambda)(x). \end{aligned}$$

We say that the BF-space B is twisted left-invariant if $\ell_y^\sigma f \in B$ for all $f \in B$ and $f \mapsto \ell_y^\sigma f$ is bounded for all $y \in G$. Analogously, we define twisted right-invariant spaces. In the sequel we will assume any BF-space to be twisted right- and left-invariant, and that twisted left and right translations by elements y in a compact set U are uniformly bounded in the sense that there is a finite constant C_U such that for any $y \in U$

$$\|\ell_y^\sigma f\| \leq C_U \|f\| \text{ and } \|r_y^\sigma f\| \leq C_U \|f\|. \quad (3.1)$$

We say that left translation, ℓ , is continuous on B if for every $f \in B$ the mapping $y \mapsto \ell_y f$ is continuous $G \rightarrow B$. Similarly for r . In general we do not assume that ℓ and r are continuous on B until we need to derive atomic decompositions in sections 5 and 6. Moreover, the twisted translations ℓ^σ and r^σ are generally not continuous (not even near e), since σ is not globally continuous. A BF-space B is called solid if $|f| \leq |g|$ and $g \in B$ implies that $f \in B$. If B is a solid space, then B is twisted left or right invariant if and only if B is left or right invariant.

Remark 3.2. A weight on G is a measurable function $w : G \rightarrow (0, \infty)$. For a weight w we define the weighted Lebesgue spaces

$$L_w^p(G) = \left\{ f \text{ measurable} \mid \|f\|_{L_w^p} := \left(\int |f(x)|^p w(x) dx \right)^{1/p} < \infty \right\}.$$

The spaces $L_w^p(G)$ are clearly solid Banach function spaces.

The weight w is called submultiplicative, if $w(xy) \leq w(x)w(y)$ for all $x, y \in G$. When w is submultiplicative the spaces $L_w^p(G)$ are left and right invariant for $1 \leq p \leq \infty$ and left and right translation are continuous for $1 \leq p < \infty$. Moreover, ℓ^σ and r^σ are projective representations on $L_w^p(G)$ for $1 \leq p < \infty$. The cocycle for ℓ^σ is σ while r^σ has cocycle $\bar{\sigma}$.

It is easily verified that if w is bounded on bounded sets, then condition (3.1) holds on $L_w^p(G)$ for $1 \leq p < \infty$. By direct inspection of the norm, it can also be verified for $p = \infty$.

We now define sequence spaces related to a solid BF-space B . Let U be a compact neighbourhood of the identity, and let $\{x_i\}$ be a countable U -dense and well-spread collection of elements in G . Define the sequence space $\dot{b}$ by

$$\dot{b} = \left\{ \{c_i\} \subseteq \mathbb{C} \mid \sum_i c_i 1_{x_i U} \in B \right\},$$

with norm

$$\|\{c_i\}\|_{\dot{b}} = \left\| \sum_i c_i 1_{x_i U} \right\|_B.$$

For example, if $B = L^p(G)$, then $\dot{b} = \ell^p$.

In the remainder of this paper we will need the following generalization of convolution in the presence of a cocycle. For a cocycle σ we define the twisted convolution of functions f and g on G by

$$f \# g(x) := \int_G f(y) \ell_y^\sigma g(x) dy = \int_G f(y) g(y^{-1}x) \overline{\sigma(y, y^{-1}x)} dy$$

whenever the integral exists. If the cocycle is constant this is the same as usual group convolution for which we reserve the special notation

$$f * g(x) := \int_G f(y) g(y^{-1}x) dy.$$

4. COORBIT SPACES

Let $u \in \mathcal{S}$ and let B be a twisted left-invariant BF-space. Define

$$\text{Co}_\rho^u B := \{\lambda \in \mathcal{S}^* \mid W_u^\rho(\lambda) \in B\}$$

with the norm

$$\|\lambda\|_{\text{Co}_\rho^u B} := \|W_u^\rho(\lambda)\|_B.$$

We will now impose conditions which ensure that $\text{Co}_\rho^u B$ is a Banach space. In the process we will demonstrate that these conditions ensure that the space

$$B_u^\# := \{f \in B \mid f \# W_u^\rho(u) = f\}$$

with norm inherited from B is a reproducing kernel Banach space isometrically isomorphic to $\text{Co}_\rho^u B$.

A ρ -cyclic vector $u \in \mathcal{S}$, is called a ρ -analyzing vector for $\mathcal{S}$ if the reproducing formula

$$W_u^\rho(\lambda) \# W_u^\rho(u) = W_u^\rho(\lambda)$$

holds for all $\lambda \in \mathcal{S}^*$.

Assumption 4.1. Let B be a twisted left-invariant BF-space on G . Assume there exists a nonzero ρ -analyzing vector $u \in \mathcal{S}$ satisfying the following continuity condition: The mapping

$$B \times \mathcal{S} \ni (f, v) \mapsto f \# W_v^\rho(u)(1) = \int_G f(y) W_v^\rho(u)^\vee(y) \overline{\sigma(y, y^{-1})} dy \in \mathbb{C}$$

is continuous.

Remark 4.2. If $B = L_w^p(G)$ then the continuity condition will be a duality requirement. Specifically, Assumption 4.1 is satisfied for $B = L_w^p(G)$ if the topology on $\mathcal{S}$ is such that

$$\mathcal{S} \ni v \mapsto W_v^\rho(u)^\vee \in L_{w^{-q/p}}^q(G)$$

is continuous, where $\frac{1}{p} + \frac{1}{q} = 1$.

The main result of this section is

Theorem 4.3. *Let $(\rho, \mathcal{S})$ be a projective representation of G , and let B be a twisted left-invariant BF-space on G . Assume that $u \in \mathcal{S}$ is a ρ -analyzing vector satisfying Assumption 4.1. Then*

- (1) $W_u^\rho(v) \# W_u^\rho(u) = W_u^\rho(v)$ for $v \in \text{Co}_\rho^u B$.
- (2) The space $\text{Co}_\rho^u B$ is a ρ^* -invariant Banach space.
- (3) $W_u^\rho : \text{Co}_\rho^u B \rightarrow B$ intertwines ρ^* and $a \mapsto \ell_a^\sigma$.
- (4) $W_u^\rho : \text{Co}_\rho^u B \rightarrow B_u^\#$ is an isometric isomorphism.

From [7] it is known that the statements are true if the cocycle σ is constant and therefore ρ is a representation. Notice that in [7] the representation ρ is assumed strongly continuous, but that requirement can be replaced by the weak continuity of $x \mapsto \langle \lambda, \rho(x)u \rangle$ for $\lambda \in \mathcal{S}^*$ and $u \in \mathcal{S}$ without modifications. We will therefore prove Theorem 4.3 by connecting it to coorbit theory for the representation ρ_σ for an appropriate choice of BF-space $\widehat{B}$ on the Mackey group. It turns out that the space

$$\widehat{B} := \{F : G \times \mathbb{T} \rightarrow \mathbb{C} \mid F(a, t) = \bar{t}f(a), f \in B\}$$

with norm $\|F\|_{\widehat{B}} := \|f\|_B$ is a good choice.

Lemma 4.4. *If $G \times \mathbb{T}$ and $\widehat{B}$ are defined as before, then the following relations hold.*

- (1) The spaces $B, \widehat{B}$ are isometrically isomorphic via $\Lambda f(x, t) := \bar{t}f(x)$.
- (2) If the space B is twisted left-invariant, then $\widehat{B}$ is left-invariant.
- (3) For $F \in \widehat{B}$, we have $F * W_u^{\rho_\sigma}(u)(x, z) = \bar{z}f \# W_u^\rho(u)(x)$ when $F(a, t) = \bar{t}f(a)$.

Proof. The first part is clear. The second part done by the following calculations:

$$\begin{aligned}
\ell_{(a,w)}F(x,z) &= F(a^{-1}x, \overline{wz\sigma(a, a^{-1})\sigma(a^{-1}, x)}) \\
&= F(a^{-1}x, \overline{wz\sigma(a, a^{-1}x)}) \\
&= w\overline{z\sigma(a, a^{-1}x)}f(a^{-1}x) \\
&= \overline{z}w\ell_a^\sigma f(x).
\end{aligned}$$

Therefore, B is twisted left-invariant if and only if $\widehat{B}$ is left invariant. For the third part, we have

$$\begin{aligned}
F * W_u(u)(x, z) &= \iint F(y, w)W_u^{\rho_\sigma}(u)((y, w)^{-1}(x, z))dw dy \\
&= \iint F(y, w)W_u^{\rho_\sigma}(u)(y^{-1}x, \overline{wz\sigma(y, y^{-1})\sigma(y^{-1}, x)})dw dy \\
&= \overline{z} \int f(y)W_u^\rho(u)(y^{-1}x) \overline{\sigma(y, y^{-1})\sigma(y^{-1}, x)} dy \\
&= \overline{z} f \# W_u^\rho(u)(x).
\end{aligned}$$

□

Lemma 4.5. *A vector u is ρ -analyzing if and only if u is ρ_σ -analyzing.*

Proof. First, we know that u is ρ_σ -cyclic if and only if u is ρ -cyclic, since $\langle \lambda, \rho(x)u \rangle = t\langle \lambda, \rho_\sigma(x, t)u \rangle$.

A straightforward calculation gives

$$W_u^{\rho_\sigma}(\lambda) * W_u^{\rho_\sigma}(u)(x, t) = \overline{t}W_u^\rho(\lambda) \# W_u^\rho(u)(x),$$

and the claim follows from this. □

The following theorem provides the connection between the coorbit theory that arises from representations [7] and the coorbit theory that arises from projective representations.

Theorem 4.6. *The triple B , ρ and u satisfy Assumption 4.1 with cocycle σ if and only if the triple $\widehat{B}$, ρ_σ and u satisfy Assumption 4.1 with a constant cocycle. Therefore $\text{Co}_\rho^u B$ and $\text{Co}_{\rho_\sigma}^u \widehat{B}$ are simultaneously defined. Moreover, $\text{Co}_\rho^u B = \text{Co}_{\rho_\sigma}^u \widehat{B}$ with the same norm.*

Proof. By Lemma 4.4 the space $\widehat{B}$ is left invariant. Next, denote the wavelet transform related to the representation ρ_σ by $W_u^{\rho_\sigma}$.

Note that if $F(x, t) = \overline{t}f(x)$, then

$$\begin{aligned}
\iint F(x, z)W_v^{\rho_\sigma}(u)((x, z)^{-1})dz dx &= \iint F(x, z)W_v^{\rho_\sigma}(u)((x^{-1}, \overline{z\sigma(x^{-1}, x)}))dz dx \\
&= \iint \overline{z}f(x)W_v^\rho(u)(x^{-1})\overline{z\sigma(x, x^{-1})}dz dx \\
&= f \# W_v^\rho(u)(1).
\end{aligned}$$

It follows that the continuity of $(f, v) \mapsto \int f(x)W_v^\rho(u)(x^{-1})\overline{\sigma(x, x^{-1})}dx$ on $B \times \mathcal{S}$ is equivalent to the continuity of $(F, v) \mapsto \iint F(x, z)W_v^{\rho_\sigma}(u)((x, z)^{-1})dz dx$.

Next, assume that $\lambda \in \mathcal{S}$. Then, by Lemma 2.12, $\lambda \in \text{Co}_\rho^u B \Leftrightarrow W_u^\rho(\lambda) \in B \Leftrightarrow W_u^{\rho_\sigma}(\lambda) \in \widehat{B} \Leftrightarrow \lambda \in \text{Co}_{\rho_\sigma}^u \widehat{B}$. □

Now we demonstrate our main result about the coorbit space constructed by the twisted convolution.

Proof of Theorem 4.3. By Theorem 4.6, the space $\widehat{B}$ and u satisfy Assumption 4.1. So we can apply Theorem 4.6 to the space $\widehat{B}$.

(1) The identity is assumed true for all analyzing vectors u and all functionals λ , and therefore it is also true for members of the coorbit space.

(2) We know that the space $\text{Co}_{\rho_\sigma}^u \widehat{B} = \text{Co}_\rho^u B$ is ρ_σ^* -invariant Banach space. So $W_u(\rho_\sigma^*(y, w)\phi) \in \text{Co}_{\rho_\sigma}^u \widehat{B}$. On the other hand

$$W_u^{\rho_\sigma}(\rho_\sigma^*(y, w)\phi)(x, z) = \overline{zw}W_u^\rho(\rho^*(y)\phi)(x),$$

which implies that $W_u^\rho(\rho^*(y)\phi) \in B$.

(3) Using the fact that $W_u^{\rho_\sigma}$ intertwines ρ_σ^* with left translation, and $\rho_\sigma^*(x, z) = z\rho^*(x)$. We have

$$\begin{aligned} W_u^\rho(\rho^*(y)\phi)(x) &= \overline{wz}W_u^{\rho_\sigma}(\rho_\sigma^*(y, w)\phi)(x, z) = \overline{wz}\ell_{(y, w)}W_u^{\rho_\sigma}(\phi)(x, z) \\ &= \overline{\sigma(y, y^{-1})}\sigma(y^{-1}, x)\ell_y W_u^\rho(\phi)(x) \\ &= \ell_y^\sigma W_u^\rho(\phi)(x). \end{aligned}$$

(4) According to (1) and (3) in Lemma 4.4 the spaces $B_u^\#$ and

$$\widehat{B}_u = \{F \in \widehat{B} \mid F * W_u^{\rho_\sigma}(u) = F\},$$

where convolution is on the group G_σ , are isometrically isomorphic. If we denote the isometrical isomorphism between B_u and $\widehat{B}_u$ by Λ , then $W_u^\rho = \Lambda^{-1}W_u^{\rho_\sigma} : \text{Co}_\rho^u B \rightarrow B_u^\#$ and the result is obtained. $\square$

In the following theorem, we prove that the twisted coorbit space is independent of the choice of the ρ -analyzing vector under some assumptions.

Theorem 4.7. *Assume that u_1 and u_2 both satisfy Assumption 4.1, and the following properties are true for $i, j \in \{1, 2\}$*

- (1) *there are nonzero constants $C_{i,j}$ such that $W_{u_i}^\rho(\lambda) \# W_{u_j}^\rho(u_i) = C_{i,j}W_{u_j}^\rho(\lambda)$ for all $\lambda \in \mathcal{S}^*$*
- (2) *the mapping $B_{u_i} \ni f \mapsto f \# W_{u_j}^\rho(u_i) \in B$ is continuous.*

Then $\text{Co}_\rho^{u_1} B = \text{Co}_\rho^{u_2} B$ with equivalent norms.

Proof. We already know from [7] that this theorem is true for representations, i.e. when $\sigma = 1$. The proof thus consists of connecting the statements for ρ with similar statements for ρ_σ .

Consider the space $\widehat{B}$ and the Mackey group $G \times \mathbb{T}$. Since u_1 and u_2 are ρ -analyzing vectors for $\mathcal{S}$ that satisfy Assumption 4.1, they are also π_ρ -analyzing vectors for $\mathcal{S}$ that satisfy Assumption 4.1 (see Theorem 4.6). Also for $i, j \in \{1, 2\}$ and $\lambda \in \mathcal{S}^*$, we have

$$\begin{aligned} W_{u_i}^{\rho_\sigma}(\lambda) * W_{u_j}^{\rho_\sigma}(u_i)(x, t) &= \overline{t}W_{u_i}^\rho(\lambda) \# W_{u_j}^\rho(u_i)(x) \\ &= \overline{t}C_{i,j}W_{u_j}^\rho(\lambda)(x) \\ &= C_{i,j}W_{u_j}^{\rho_\sigma}(\lambda)(x, t) \end{aligned}$$

Moreover, the mapping $\widehat{B}_{u_i} \ni F \mapsto F * W_{u_j}^{\rho_\sigma}(u_i) \in \widehat{B}$ is continuous, indeed,

$$\|F * W_{u_j}^{\rho_\sigma}(u_i)\|_{\widehat{B}} = \|f \# W_{u_j}^\rho(u_i)\|_B \leq C\|f\|_B = C\|F\|_{\widehat{B}}.$$

Therefore, by Theorem 2.7 in [7], $\text{Co}_{\rho\sigma}^{u_1}\widehat{B} = \text{Co}_{\rho\sigma}^{u_2}\widehat{B}$. Since $\text{Co}_{\rho\sigma}^{u_i}\widehat{B} = \text{Co}_{\rho}^{u_i}B$, the result is obtained. $\square$

5. ATOMIC DECOMPOSITIONS AND FRAMES

The atomic decompositions we provide build on sampling on the reproducing kernel space $B_u^\#$. To do so, we need to show that the reproducing kernel does not vary too much under translation by elements of a fixed compact neighbourhood. The special form of the reproducing kernel allow us to estimate such local variations using smoothness of the analyzing vector u .

Definition 5.1. A vector $u \in \mathcal{S}$ is called ρ -weakly differentiable if for all $X \in \mathfrak{g}$ and for all $\lambda \in \mathcal{S}^*$ the mapping $t \mapsto \langle \lambda, \rho(\exp(tX))u \rangle$ is differentiable. In this case we denote by $\rho(X)u$ the vector in $\mathcal{S}$ satisfying

$$\langle \lambda, \rho(X)u \rangle = \left. \frac{d}{dt} \right|_{t=0} \langle \lambda, \rho(\exp(tX))u \rangle.$$

Similarly, a vector $\lambda \in \mathcal{S}$ is called ρ^* -weakly differentiable if for all $X \in \mathfrak{g}$ and for all $u \in \mathcal{S}$ the mapping $t \mapsto \langle \rho^*(\exp(tX))\lambda, u \rangle$ is differentiable. In this case we denote by $\rho^*(X)\lambda$ the distribution in $\mathcal{S}^*$ satisfying

$$\langle \rho^*(X)\lambda, u \rangle = \left. \frac{d}{dt} \right|_{t=0} \langle \rho^*(\exp(tX))\lambda, u \rangle.$$

Assumption 5.2. Let $(\rho, \mathcal{S})$ be a projective representation of G , and let B be a solid BF-space on G on which left and right translation are continuous. Assume that B , ρ and u satisfies Assumption 4.1. Assume, moreover, that u is ρ -weakly and ρ^* -weakly differentiable up to order $n = \dim(G)$, and that for all finite subsets with n elements, $\{Y_1, Y_2, \dots, Y_n\} \subseteq \mathfrak{g}$ the mappings

$$f \mapsto f * |W_u^\rho(\rho^*(Y_1)\rho^*(Y_2) \cdots \rho^*(Y_n)u)| \text{ and } f \mapsto f * |W_{\rho(Y_1)\rho(Y_2) \cdots \rho(Y_n)u}^\rho(u)|$$

are bounded on the solid, left and right invariant BF-space B .

Theorem 5.3. Let u be a vector satisfying Assumption 5.2. Let $X_1, \dots, X_n$ be a fixed basis for $\mathfrak{g}$ and define $U_\epsilon = \{e^{t_1 X_1} \cdots e^{t_n X_n} \mid -\epsilon \leq t_k \leq \epsilon\}$. In the following we always choose a cocycle σ and $\epsilon > 0$ small enough that σ is C^∞ on a neighbourhood containing $U_\epsilon \times U_\epsilon$.

- (1) Given a U_ϵ -dense and well-spread sequence $\{x_i\} \subseteq G$ and an U_ϵ -BUPU $\{\psi_i\}$, then the operators

$$\begin{aligned} T_1 f &= \sum_i f(x_i) \sigma(x, x^{-1}x_i) \psi_i \# W_u^\rho(u) \\ T_2 f &= \sum_i \lambda_i(f) \ell_{x_i}^\sigma W_u^\rho(u) \\ T_3 f &= \sum_i c_i f(x_i) \ell_{x_i}^\sigma W_u^\rho(u) \end{aligned}$$

where $\lambda_i(f) = \int f(y) \psi_i(y) \overline{\sigma(y, y^{-1}x_i)} dy$ and $c_i = \int \psi_i(y) dy$, are well defined on B_u^σ .

- (2) There is an ϵ small enough that the operators T_1, T_2, T_3 are invertible for any U_ϵ -dense and well-spread sequence $\{x_i\} \subseteq G$ and any U_ϵ -BUPU $\{\psi_i\}$. In this case the family $\{\rho^*(x_i)u\}$ is a Banach frame for $\text{Co}_\rho^u B$ with respect to the sequence space $\dot{b}$, and the families $\{\lambda_i \circ T_2^{-1} \circ W_u^\rho, \rho^*(x_i)u\}$ and

$\{c_i T_3^{-1} \circ W_u^\rho, \rho^*(x_i)u\}$ are atomic decompositions for $\text{Co}_\rho^u B$ with respect to the sequence space $\dot{b}$. In particular, $\gamma \in \text{Co}_\rho^u B$ can be reconstructed by

$$\gamma = (W_u^\rho)^{-1} T_1^{-1} \left(\sum_i W_u^\rho(\gamma)(x_i) \psi_i \# W_u^\rho(u) \right)$$

$$\gamma = \sum_i \lambda_i (T_2^{-1} W_u^\rho(\gamma)) \rho^*(x_i) u$$

$$\gamma = \sum_i c_i (T_3^{-1} W_u^\rho(\gamma)) \rho^*(x_i) u$$

with convergence in $\mathcal{S}^*$. The convergence is in $\text{Co}_\rho^u B$ if $C_c(G)$ is dense in B .

These results can be proven by applying Theorems 3.4, 3.6 and 3.7 from [5] to the representation ρ_σ of G_σ . To do so we need to define an appropriate solid BF-space on G_σ . There are many choices for such BF-spaces and two different approaches have been presented in [9] and [16]. However, we find it useful to avoid making this choice and to work directly on the space B . The calculations differ non-trivially from [5] by the occurrence of the cocycle, and the details are carried out in the appendix. It should also be mentioned that the results in the appendix do allow for the use of non-differentiable u in case there is another way to estimate the local oscillations of the kernel.

6. BERGMAN SPACES ON THE UNIT BALL

Recently, in [4], the first and the third authors together with Gröchenig, obtained atomic decompositions for Bergman spaces on the unit ball in $\mathbb{C}^n$ through a finite covering group of the group $\text{SU}(n, 1)$ with the restriction that the representation parameter $s > n$ had to be rational. As a special case, atomic decompositions of Bergman spaces through the group $\text{SU}(n, 1)$ is valid for integer values of the parameter $s > n$. Overcoming the restriction on the parameter is one reason for introducing a coorbit theory for projective representations.

We dedicate this chapter to generating Banach frames and atomic decompositions of Bergman spaces on the unit ball via the group $\text{SU}(n, 1)$. For more references we encourage the reader to see [3, 17, 23, 24, 32, 34].

6.1. Facts about Bergman Spaces on the Unit Ball. In this section we collect facts about Bergman spaces on the unit ball. Let $\mathbb{C}^n$ be equipped with the usual inner product $(z, w) = z_1 \overline{w_1} + z_2 \overline{w_2} + \dots + z_n \overline{w_n}$ and define the unit ball by

$$\mathbb{B}^n := \{z \in \mathbb{C}^n \mid |z|^2 := |z_1|^2 + |z_2|^2 + \dots + |z_n|^2 < 1\}.$$

Let dv be a normalized volume measure on the unit ball under identifying $\mathbb{C}^n$ with $\mathbb{R}^{2n}$. For $\alpha > -1$, define the measure $dv_\alpha(z) := C_\alpha (1 - |z|^2)^\alpha dv(z)$ where C_α is a constant such that dv_α is a probability measure. Notice that the measure dv_α is finite measure on $\mathbb{B}^n$ if and only if $\alpha > -1$.

We define the α -weighted L^p space on the unit ball as

$$L_\alpha^p(\mathbb{B}^n) = \{f : \mathbb{B}^n \rightarrow \mathbb{C} \mid \int_{\mathbb{B}^n} |f(z)|^p dv_\alpha(z) < \infty\}$$

with norm

$$\|f\|_{L_\alpha^p} = \left(\int_{\mathbb{B}^n} |f(z)|^p dv_\alpha(z) \right)^{1/p},$$

where $1 \leq p < \infty$. For $\alpha > -1$, we define the weighted Bergman spaces on the unit ball to be

$$A_\alpha^p(\mathbb{B}^n) := L_\alpha^p(\mathbb{B}^n) \cap \mathcal{O}(\mathbb{B}^n)$$

with norm inherited from $L_\alpha^p(\mathbb{B}^n)$, where $\mathcal{O}(\mathbb{B}^n)$ is the space of holomorphic functions on the unit ball. We have the condition $\alpha > -1$ to construct a non-trivial Bergman spaces, in fact, if $\alpha \leq -1$, then the only holomorphic function in $L_\alpha^p(\mathbb{B}^n)$ is the zero function.

As we have seen in the special case on the unit disc, Bergman spaces are closed subspaces of $L_\alpha^p(\mathbb{B}^n)$, *i.e.*, Bergman spaces are Banach spaces. In the case $p = 2$, the space $A_\alpha^2(\mathbb{B}^n)$ is a Hilbert space with the inner product

$$(f, g)_\alpha = \int_{\mathbb{B}^n} f(z) \overline{g(z)} dv_\alpha(z).$$

The orthogonal projection of $L_\alpha^2(\mathbb{B}^n)$ on the space $A_\alpha^2(\mathbb{B}^n)$ is given by

$$P_\alpha f(z) = \int_{\mathbb{B}^n} f(w) K_\alpha(z, w) dv_\alpha(w),$$

where

$$K_\alpha(z, w) = \frac{1}{(1 - (z, w))^{n+1+\alpha}}$$

is the reproducing kernel for $A_\alpha^2(\mathbb{B}^n)$.

The group $\mathrm{SU}(n, 1)$ is defined to be the group of all $(n+1) \times (n+1)$ -matrices x of determinant 1 and $x^* J_{(n,1)} x = J_{(n,1)}$, where

$$J_{(n,1)} = \begin{pmatrix} -I_n & 0 \\ 0 & 1 \end{pmatrix}$$

We always write $x \in \mathrm{SU}(n, 1)$ in the block form

$$x = \begin{pmatrix} A & b \\ c^t & d \end{pmatrix},$$

where A is an $n \times n$ matrix, and b, c are vectors in $\mathbb{C}^n$, and $d \in \mathbb{C}$. Simple calculations show that

$$x^{-1} = \begin{pmatrix} A^* & -\bar{c} \\ -\bar{b}^t & \bar{d} \end{pmatrix}.$$

The fact that $xx^{-1} = I$ implies

$$|d|^2 - |b|^2 = 1 \tag{6.1}$$

Form now on, we write $G = \mathrm{SU}(n, 1)$. This group acts transitively on $\mathbb{B}^n$ by

$$x \cdot z = (Az + b)((c, \bar{z}) + d)^{-1}.$$

If we define the subgroup K of G as

$$K = \left\{ \begin{pmatrix} k & 0 \\ 0 & \frac{1}{\det(k)} \end{pmatrix} \mid k \in U(n) \right\},$$

then the stabilizer of the origin $o \in \mathbb{C}^n$ is K and $\mathbb{B}^n \simeq G/K$. It follows that there is a one to one correspondence between the K -right invariant functions on G and the functions on $\mathbb{B}^n$ via

$$\tilde{f}(x) = f(x \cdot 0).$$

This correspondence defines an G -invariant measure on $\mathbb{B}^n$ which is given by $(1 - |z|^2)^{-n-1} dz$. The compactness of K ensures that we can normalize this measure so that, for any $\tilde{f}$ K -right invariant function on G , we have

$$\int_G \tilde{f}(x) dx = \int_{\mathbb{B}^n} f(z) (1 - |z|^2)^{-n-1} dz. \quad (6.2)$$

Let $v_\alpha(x) = (1 - |x \cdot o|^2)^\alpha$, then we define the v_α -weighted L^p spaces on G to be

$$L_\alpha^p(G) = \left\{ F : G \rightarrow \mathbb{C} \left| \|F\|_{L_\alpha^p(G)} := \left(c_\alpha \int_G |F(x)|^p (1 - |x \cdot o|^2)^\alpha dx \right)^{1/p} < \infty \right. \right\}$$

If we denote by $L_\alpha^p(G)^K$ the space of K -right invariant functions in the space $L_\alpha^p(G)$, then it is easy to see that $L_\alpha^p(\mathbb{B}^n)$ and $L_{\alpha+n+1}^p(G)^K$ are isometric. That is,

$$\|f\|_{L_\alpha^p(\mathbb{B}^n)} = \|\tilde{f}\|_{L_{\alpha+n+1}^p(G)}. \quad (6.3)$$

For $s > n$, the action of G on $\mathbb{B}^n$ defines an irreducible unitary projective representation of G on the space $\mathcal{H}_s = A_{s-n-1}^2$ by

$$\rho_s(x)f(z) = (-(z, b) + \bar{d})^{-s} f(x^{-1} \cdot z), \quad (6.4)$$

which also defines a representation for the universal covering group of G . From now on we assume that a cocycle σ has been chosen for the projective representation ρ_s . Note, that since ρ_s is a local representation, the cocycle can be chosen equal to one on a neighbourhood of $e \times e$ (see also Theorem 7.1 in [2]).

We denote the twisted wavelet transform on $\mathcal{H}_s$ by

$$W_u^{\rho_s}(\lambda)(x) = (\lambda, \rho_s(x)u)_{\mathcal{H}_s}.$$

Let $\mathcal{P}_k$ be the space of all homogeneous polynomials of degree k on $\mathbb{C}^n$. In the following theorem we summarize some properties of the space of smooth vectors for ρ_s and its conjugate dual space, which will be the candidate Fréchet space $\mathcal{S}$ for constructing the coorbits of $L_{\alpha+n+1-sp/2}^p(G)$.

Theorem 6.1. *Let $s > n$ and let $(\rho_s, \mathcal{H}_s)$ be the projective representation of G which is defined in 6.4. The following are true:*

- (1) *Every polynomial is a smooth vector for ρ_s .*
- (2) *Every smooth vector for ρ_s is bounded.*
- (3) *Assume $v \in \mathcal{H}_s$, then $v \in \mathcal{H}_s^\infty$ if and only if $v = \sum_k v_k$, $v_k \in \mathcal{P}_k$, and for all $N \in \mathbb{N}$ there exists a constant $C_N > 0$ such that $\|v_k\|_{\mathcal{H}_s} \leq C_N(1+k)^{-N}$.*
- (4) *A vector $\phi \in \mathcal{H}_s^{-\infty}$ if and only if $\phi = \sum_k \phi_k$, $\phi_k \in \mathcal{P}_k$, and there exist $N \in \mathbb{N}$ and $C > 0$ such that $\|\phi_k\|_{\mathcal{H}_s} \leq C(1+k)^N$. Moreover, the dual pairing is given by*

$$\langle \phi, v \rangle_s = \sum_k (\phi_k, v_k)_{\mathcal{H}_s}.$$

Proof. The proof is done by noting that ρ_s is a unitary representation of the universal covering group of G , so the smooth vectors are the same for both, where the smooth vectors for ρ_s , as a representation, satisfy all the above properties as proved in [3] and [4]. $\square$

6.2. Bergman Spaces as Coorbits spaces. In this section we collect several facts from [4] and use them with some modifications to provide atomic decompositions of Bergman spaces through projective representations of the group $SU(n, 1)$.

As before, we assume $G = SU(n, 1)$ and $(\mathcal{H}_s^\infty, \rho_s)$ is the smooth projective representation obtained by restricting $(\mathcal{H}_s, \rho_s)$ to $\mathcal{H}_s^\infty$. In this section we show that Bergman spaces are twisted convolutive coorbits of weighted L^p spaces, which allows us to discretize Bergman spaces using the full group $SU(n, 1)$. For this goal we need the following results which already proved for the linear representation in [4]. The same proof will work (with minor differences) for the projective representation case. For completeness we will provide a full proof for each of these results.

Lemma 6.2. *Assume u and v are smooth vectors for ρ_s . There is a constant C depending on u and v such that*

$$|W_u^{\rho_s}(v)(x)| \leq C(1 - |x \cdot o|^2)^{s/2} (1 - \log(1 - |x \cdot o|^2)).$$

Proof. Let $x = \begin{pmatrix} A & b \\ c^t & d \end{pmatrix}$. Since smooth vectors for ρ_s are bounded (see Theorem 6.1), we can define $C = \sup_{z, w \in \mathbb{B}^n} |u(z)||v(w)|$. Now, note that

$$\begin{aligned} |\rho_s(x)u(z)| &= |\bar{d} - (z, b)|^{-s} |u(x^{-1} \cdot z)| \\ &= |d|^{-s} |1 - (z, bd^{-1})|^{-s} |u(x^{-1} \cdot z)| \\ &= |d|^{-s} |1 - (z, x \cdot 0)|^{-s} |u(x^{-1} \cdot z)|. \end{aligned}$$

Also, by (6.1) we have $|d|^{-2} = 1 - |bd^{-1}|^2 = 1 - |x \cdot o|^2$. In other words, $|d|^{-s} = (1 - |x \cdot o|^2)^{s/2}$. It follows that

$$\begin{aligned} |W_u^{\rho_s}(v)(x)| &= \left| \int_{\mathbb{B}^n} v(z) \overline{\rho_s(x)u(z)} (1 - |z|^2)^{s-n-1} dz \right| \\ &\leq |d|^{-s} \int_{\mathbb{B}^n} |v(z)| |u(x^{-1} \cdot z)| |1 - (z, x \cdot 0)|^{-s} (1 - |z|^2)^{s-n-1} dz \\ &\leq C(1 - |x \cdot o|^2)^{s/2} \int_{\mathbb{B}^n} |1 - (z, x \cdot 0)|^{-s} (1 - |z|^2)^{s-n-1} dz. \end{aligned}$$

The last integral is comparable to $1 - \log(1 - |x \cdot o|^2)$ (see [34, Theorem 1.12]), which proves our assertion. $\square$

Proposition 6.3. *Let $\alpha > -1$, $1 \leq p < \infty$, and $s > n$ be chosen. Assume that u and v are smooth vectors for ρ_s . Then $W_u^{\rho_s}(v) \in L_t^p(G)$ for $t + ps/2 > n$.*

Proof. Assume that $u, v \in \mathcal{H}_s^\infty$. Since for any $\epsilon > 0$ the limit

$$\lim_{t \rightarrow 1} (1 - t)^\epsilon (1 - \log(1 - t)) = 0$$

where $t \in (0, 1)$, one can find $C_1 > 0$ such that

$$1 - \log(1 - |x \cdot o|^2) \leq C_1(1 - |x \cdot o|^2)^{-\epsilon/2}$$

for all $x \in G$. Now, for $t + ps/2 > n$ choose ϵ small enough so that $t + p(s - \epsilon)/2 > n$. By Lemma 6.2, there is a constant C_2 such that

$$|W_u^{\rho_s}(v)(x)| \leq C_2(1 - |x \cdot o|^2)^{s/2} (1 - \log(1 - |x \cdot o|^2)) \leq C(1 - |x \cdot o|^2)^{(s-\epsilon)/2}.$$

Since the function $x \mapsto (1 - |x \cdot o|^2)$ is K -right invariant on G , we can write the following integral as an integral on $\mathbb{B}^n$ as in (6.2)

$$\begin{aligned} \int_G |W_u^{\rho_s}(v)(x)|^p (1 - |x \cdot o|)^t dx &\leq C \int_G (1 - |x \cdot o|^2)^{(s-\epsilon)p/2+t} dx \\ &= C \int_{\mathbb{B}^n} (1 - |z|^2)^{(s-\epsilon)p/2+t-n-1} dz. \end{aligned}$$

For $t + sp/2 > t + (s - \epsilon)p/2 > n$, we have $(s - \epsilon)p/2 + t - n - 1 > -1$. Therefore, the last integral is finite, and hence, $W_u^{\rho_s}(v) \in L_t^p(G)$ for $t + sp/2 > n$. $\square$

Now we show that the twisted coorbits of the spaces $L_{\alpha+n+1-sp/2}^p(G)$ generated by any nonzero smooth vector $u \in \mathcal{H}_s^\infty$ are well defined nonzero spaces under the assumptions in the following theorem.

Theorem 6.4. *Let $1 \leq p < \infty$, and $s > n$. Assume that $-1 < \alpha < p(s - n) - 1$. For a nonzero smooth vector $u \in \mathcal{H}_s^\infty$, the coorbit space $\text{Co}_{\rho_s}^u L_{\alpha+n+1-sp/2}^p(G)$ is a nonzero well defined Banach space.*

Proof. Let us show that the nonzero smooth vector $v \in \mathcal{H}_s^\infty$ satisfies Assumption 4.1. First, u is ρ -cyclic because $\mathcal{H}_s^\infty$ is irreducible projective representation. Next, since ρ_s is square integrable and G is unimodular, every smooth vector for ρ_s is ρ -admissible, i.e., u is in the domain of the operator A_ρ , which is given in Theorem 2.9. It follows that

$$\begin{aligned} W_u^{\rho_s}(v) \# W_u^{\rho_s}(u)(x) &= \int_G (v, \rho(y)u)_{\mathcal{H}_s} (u, \rho(y^{-1}x)u)_{\mathcal{H}_s} \overline{\sigma(y, y^{-1}x)} dy \\ &= \int_G (v, \rho(y)u)_{\mathcal{H}_s} (u, \rho(y^{-1})\rho(x)u)_{\mathcal{H}_s} \overline{\sigma(y^{-1}, x)} \overline{\sigma(y, y^{-1}x)} dy \\ &= \int_G (v, \rho(y)u)_{\mathcal{H}_s} (u, \rho(y)^{-1}\rho(x)u)_{\mathcal{H}_s} \sigma(y, y^{-1}) \overline{\sigma(y, y^{-1})} dy \\ &= \int_G (v, \rho(y)u)_{\mathcal{H}_s} (\rho(y)u, \rho(x)u)_{\mathcal{H}_s} dy \\ &= c_\rho^2 (v, \rho(x)u)_{\mathcal{H}_s} (u, u)_{\mathcal{H}_s} \\ &= C W_u^{\rho_s}(v)(x) \end{aligned}$$

for all $v \in \mathcal{H}_s^\infty$. Let us show that this formula extends for all $\phi \in \mathcal{H}_s^{-\infty}$. Assume that $v \in \mathcal{H}_s^\infty$. By the orthogonality relation in Theorem 2.9, we have

$$\int_G (v, \rho(x)u)_{\mathcal{H}_s} (\rho(x)u, u)_{\mathcal{H}_s} dx = c_\rho^2 \|u\|_{\mathcal{H}_s} (v, u)_{\mathcal{H}_s}.$$

If we define $\eta := c_\rho^2 \|u\|_{\mathcal{H}} v$, then $\eta \in \mathcal{H}_s^\infty$, and hence $\langle \phi, \eta \rangle_s$ is well defined for all $\phi \in \mathcal{H}_s^{-\infty}$. By Theorem 6.1, we have

$$\begin{aligned} \langle \phi, \eta \rangle_s &= \sum_k (\phi_k, \eta)_{\mathcal{H}_s} \\ &= \sum_k c_\rho^2 \|u\|_{\mathcal{H}} (\phi_k, u)_{\mathcal{H}_s} \\ &= \sum_k \int_G (\phi_k, \rho(x)u)_{\mathcal{H}_s} (\rho(x)u, u)_{\mathcal{H}_s} dx \\ &= \int_G (\phi, \rho(x)u)_{\mathcal{H}_s} (\rho(x)u, u)_{\mathcal{H}_s} dx. \end{aligned}$$

The interchanging of the integral and the sum is valid by Tonelli's theorem, because

$$\sum_k \int_G (\phi_k, \rho(x)u)_{\mathcal{H}_s} (\rho(x)u, u)_{\mathcal{H}_s} dx = \langle \phi, \eta \rangle_s$$

exists. Therefore, the mapping

$$\phi \mapsto \int_G (\phi, \rho(x)u)_{\mathcal{H}_s} (\rho(x)u, u)_{\mathcal{H}_s} dx$$

is weakly continuous on $\mathcal{H}_s^{-\infty}$. Hence, the reproducing formula extends for all $\phi \in \mathcal{H}_s^{-\infty}$. This shows that u is a ρ -analyzing vector.

Now, we show that the mapping $(f, v) \mapsto \int_G f(x) W_u^{\rho_s}(x^{-1}) \overline{\sigma(x, x^{-1})} dx$ is continuous on $L_{\alpha+n+1-sp/2}^P(G)$. By Remark 4.2, it is enough to show that

$$W_u^{\rho_s}(v) \in (L_{\alpha+n+1-sp/2}^P(G))^* = L_{sq/2-(\alpha+n+1)q/p}^q(G),$$

where $1/p + 1/q = 1$. This is done by Proposition 6.3, because $(sq/2 - (\alpha + n + 1)q/p) + sp/2 > n$, whenever $-1 < \alpha < p(s - n) - 1$. Therefore, the space $\text{Co}_{\rho_s}^u L_{\alpha+n+1-sp/2}^P(G)$ is well defined. Finally, note that $W_u^{\rho_s}(u) \in L_{\alpha+n+1-sp/2}^P(G)$ again by Proposition 6.3, hence it is nonzero Banach space. $\square$

Our goal now is to describe Bergman spaces as twisted coorbits generated by any nonzero smooth vector $u \in \mathcal{H}_s^\infty$. First we describe Bergman spaces as twisted coorbits by the special ρ -analyzing vector $u = 1_{\mathbb{B}^n}$, then we show that this coorbit is independent of the choice of u .

Theorem 6.5. *Let $\alpha > -1$, $1 \leq p < \infty$, and $u = 1_{\mathbb{B}^n}$. The Bergman space $A_\alpha^p(\mathbb{B}^n)$ is the twisted coorbit space of $L_{\alpha+n+1-sp/2}^P(G)$ that corresponds to the projective representation $(\mathcal{H}_s^\infty, \rho_s)$. i.e., $A_\alpha^p(\mathbb{B}^n) = \text{Co}_{\rho_s}^u L_{\alpha+n+1-sp/2}^P(G)$ for $\alpha < p(s - n) - 1$.*

Proof. As in [4, Theorem 3.6], the space $A_\alpha^p(\mathbb{B}^n) \subset \mathcal{H}_s^{-\infty}$ for all $p \geq 1$, which is still valid in the case of smooth vectors for ρ . The reason behind that is that the smooth vectors for the projective representation ρ_s are the same smooth vectors of the representation of the universal covering of G . So we have only to show that for a holomorphic function f , the function $f \in L_\alpha^p(\mathbb{B}^n)$ if and only if $W_u^{\rho_s}(f) \in$

$L_{\alpha+n+1-sp/2}^p(G)$. To this end, assume $x = \begin{pmatrix} A & b \\ c^t & d \end{pmatrix}$. Then we have

$$\begin{aligned} |W_u^{\rho_s}(f)(x)| &= \left| \sum_k W_u^{\rho_s}(f_k)(x) \right| \\ &= \left| \sum_k (\bar{d})^{-s} f(bd^{-1}) \right| \\ &= |d|^{-s} \left| \sum_k f_k(x \cdot o) \right| \\ &= |d|^{-s} |f(x \cdot o)|. \end{aligned}$$

As we have seen before, $|d|^{-s} = (1 - |x \cdot o|^2)^{s/2}$. It follows that

$$|f(x \cdot o)| = (1 - |x \cdot o|^2)^{-s/2} |W_u^{\rho_s}(f)(x)|$$

by the isometry in (6.3). We conclude that $f \in L_{\alpha}^p(\mathbb{B}^n)$ if and only if $W_u^{\rho_s}(f) \in L_{\alpha+n+1-sp/2}^p(G)$. $\square$

To prove our main result in this section, which says that Bergman spaces are twisted coorbits for weighted L^p spaces generated by any smooth vector, we need the following theorem. It will be used in the subsequent section to generate a Banach frame and atomic decomposition for Bergman spaces.

Theorem 6.6. *Let $1 \leq p < \infty$, $-1 < \alpha < p(s - n) - 1$, and let v and u be smooth vectors. The convolution operator $f \mapsto f * |W_u^{\rho_s}(v)|$ is continuous on $L_{\alpha+n+1-sp/2}^p(G)$. In particular, $f \mapsto f \# W_u^{\rho_s}(v)$ is continuous on $L_{\alpha+n+1-sp/2}^p(G)$.*

Proof. Let $F \in L_{\alpha+n+1-sp/2}^p(G)$ and define

$$\tilde{f}(x) := \int_K F(xk) dk.$$

Then $\tilde{f}$ is K -right invariant function on G . Therefore, there is a corresponding $f \in L_{\alpha-sp/2}^p(\mathbb{B}^n)$. Now, for ϵ small enough such that $-(s - \epsilon)p/2 < \alpha - sp/2 + 1 < p((s - \epsilon)/2 - n) - 1$ whenever $-sp/2 < \alpha - sp/2 + 1 < p(s/2 - n) - 1$, we have

$$\begin{aligned} |F| * |W_u^{\rho_s}(v)|(x) &= \int_G |F(y)| |W_u^{\rho_s}(v)(y^{-1}x)| dy \\ &\leq C \int_G |F(y)| (1 - |y^{-1}x \cdot o|^2)^{s/2} |1 - \log(1 - |y^{-1}x \cdot o|^2)| dy \\ &\leq C \int_G |F(y)| (1 - |y^{-1}x \cdot o|^2)^{(s-\epsilon)/2} dy \\ &= C \int_{G/K} |\tilde{f}(y)| (1 - |y^{-1}x \cdot o|^2)^{(s-\epsilon)/2} dy. \end{aligned}$$

If we assume that $x = \begin{pmatrix} A_x & b_x \\ c_x^t & d_x \end{pmatrix}$, $y = \begin{pmatrix} A_y & b_y \\ c_y^t & d_y \end{pmatrix}$, $w = x \cdot o = b_x d_x^{-1}$, and $z = y \cdot o = b_y d_y^{-1}$, then

$$d_{y^{-1}x} = \bar{d}_y d_x (1 - (w, z))$$

and

$$|d_x|^{-(s-\epsilon)} = (1 - |x \cdot o|^2)^{(s-\epsilon)/2}.$$

Therefore,

$$\begin{aligned} (1 - |y^{-1}x \cdot o|^2)^{(s-\epsilon)/2} &= |d_{y^{-1}x}|^{-(s-\epsilon)} \\ &= (1 - |x \cdot o|^2)^{(s-\epsilon)/2} (1 - |y \cdot o|^2)^{(s-\epsilon)/2} (1 - (x \cdot o, y \cdot o))^{-(s-\epsilon)}. \end{aligned}$$

Thus,

$$\begin{aligned} |F| * |W_u^{\rho_s}(v)|(x) &= C \int_{G/K} |\tilde{f}(y)| \frac{(1 - |x \cdot o|^2)^{(s-\epsilon)/2} (1 - |y \cdot o|^2)^{(s-\epsilon)/2}}{|1 - (x \cdot o, y \cdot o)|^{(s-\epsilon)}} dy \\ &= C(1 - |w|^2)^{(s-\epsilon)/2} \int_{\mathbb{B}^n} |f(z)| \frac{(1 - |z|^2)^{(s-\epsilon)/2 - n - 1}}{|1 - (w, z)|^{(s-\epsilon)}} dz. \end{aligned}$$

According to [34, Theorem 2.10], the operator S which is given by

$$Sf(z) = (1 - |w|^2)^{(s-\epsilon)/2} \int_{\mathbb{B}^n} |f(z)| \frac{(1 - |z|^2)^{(s-\epsilon)/2 - n - 1}}{|1 - (w, z)|^{(s-\epsilon)}} dz$$

is continuous on $L_{\alpha-sp/2}^p(\mathbb{B}^n)$ whenever

$$-(s - \epsilon)p/2 < \alpha - sp/2 + 1 < p((s - \epsilon)/2 - n) - 1,$$

which is equivalent to $-1 < \alpha < p(s - n) - 1$. Since

$$\|f\|_{L_{\alpha-sp/2}^p(\mathbb{B}^n)} = \|\tilde{f}\|_{L_{\alpha+n+1-sp/2}^p(G/K)} = \|F\|_{L_{\alpha+n+1-sp/2}^p(G)},$$

the operator $F \mapsto F * |W_u^{\rho_s}(v)|$ is continuous on $L_{\alpha+n+1-sp/2}^p(G)$. The second part is clear from the relation $|F \# W_u^{\rho_s}(v)(x)| \leq |F| * |W_u^{\rho_s}(v)(x)|$. $\square$

We conclude our section with the following main result.

Theorem 6.7. *Let $1 \leq p < \infty$ and $-1 < \alpha < p(s - n) - 1$, and let $v \in \mathcal{H}_s^\infty$ be a nonzero smooth vector. The Bergman space $A_\alpha^p(\mathbb{B}^n)$ is the twisted coorbit space of $L_{\alpha+n+1-sp/2}^p(G)$ via the projective representation $(\mathcal{H}_s^\infty, \rho_s)$. That is, $A_\alpha^p(\mathbb{B}^n) = \text{Co}_{\rho_s}^v L_{\alpha+n+1-sp/2}^p(G)$ for $\alpha < p(s - n) - 1$.*

Proof. Assume $u = 1_{\mathbb{B}^n}$. By Theorem 6.6, we have $A_\alpha^p(\mathbb{B}^n) = \text{Co}_\rho^u L_{\alpha+n+1-sp/2}^p(G)$. We will show that the twisted coorbit $\text{Co}_{\rho_s}^v L_{\alpha+n+1-sp/2}^p(G)$ does not depend on the analyzing vector v , by applying Theorem 4.7. First, according to Theorem 6.6, the operators $f \mapsto f \# W_u^{\rho_s}(v)$ and $f \mapsto f \# W_v^{\rho_s}(u)$ are continuous on $L_{\alpha+n+1-sp/2}^p(G)$. Next, we show that $W_u^{\rho_s}(\phi) \# W_v^{\rho_s}(u) = CW_v^{\rho_s}(\phi)$ for all $\phi \in \mathcal{H}_s^\infty$. For $f \in \mathcal{H}_s^\infty$, we can use the orthogonality relation in Theorem 2.9 to get $W_u^{\rho_s}(f) \# W_v^{\rho_s}(u) = CW_v^{\rho_s}(f)$. To extend this relation to the dual of the smooth vectors, it is enough to show that

$$\phi \mapsto \int_G \langle \phi, \rho(x)u \rangle \langle \rho(x)v, u \rangle dx$$

is weakly continuous. Same argument, as in the proof of Theorem 6.4, can be made to show our claim. Therefore, the twisted coorbit spaces $\text{Co}_{\rho_s}^v L_{\alpha+n+1-sp/2}^p(G)$ are all equal to the space $A_\alpha^p(\mathbb{B}^n)$. $\square$

6.3. New atomic decompositions and frame for Bergman Spaces. In this section we generate a wavelet frame and an atomic decomposition of Bergman spaces depending on the coorbit theory, where this discretization would work for all projective representations with $s > n$, including the non-integrable cases. Also, we have more freedom in choosing the wavelet u . That is we show that any nonzero smooth vector is a good candidate to generate a Banach frame and an atomic decomposition for Bergman spaces.

Theorem 6.8. *Assume that $1 \leq p < \infty$, $s > n$, and $-1 < \alpha < p(s - n) - 1$. For a nonzero smooth vector u for ρ_s , we can choose ϵ small enough such that for every U_ϵ -well spread set $\{x_i\}_{i \in I}$ in G the following hold.*

- (1) *(Twisted wavelet frame) The family $\{\rho_s(x_i)u : i \in I\}$ is a Banach frame for $A_\alpha^p(\mathbb{B}^n)$ with respect to the sequence space $\ell_{\alpha+n+1-ps/2}^p(I)$. That is, there exist constants $A, B > 0$ such that for all $f \in A_\alpha^p(\mathbb{B}^n)$ we have*

$$A\|f\|_{A_\alpha^p(\mathbb{B}^n)} \leq \|\langle f, \rho_s(x_i)u \rangle\|_{\ell_{\alpha+n+1-ps/2}^p(I)} \leq B\|f\|_{A_\alpha^p(\mathbb{B}^n)},$$

and f can be reconstructed by

$$f = (W_u^{\rho_s})^{-1} S_1^{-1} \left(\sum_i W_u^{\rho_s}(f)(x_i) \psi_i \# W_u^{\rho_s}(u) \right)$$

where $\{\psi_i\}$ is any U_ϵ -BUPU with $\text{supp} \psi_i \subset x_i U_\epsilon$.

- (2) *(Atomic decomposition) There exists a family of functionals $\{\gamma_i\}_{i \in I}$ on $A_\alpha^p(\mathbb{B}^n)$ such that the family $\{\gamma_i, \rho_s(x_i)u\}$ forms an atomic decomposition for $A_\alpha^p(\mathbb{B}^n)$ with respect to the sequence space $\ell_{\alpha+n+1-ps/2}^p(I)$, so that any $f \in A_\alpha^p(\mathbb{B}^n)$ can be reconstructed by*

$$f = \sum_i \gamma_i(f) \rho_s(x_i)u.$$

Proof. We show that the assumptions of Theorem 5.3 are satisfied. Under the conditions on p and s , the twisted coorbit of $L_{\alpha+n+1-ps/2}^p(G)$ is well defined and u satisfies Assumption 4.1 as we have seen in Theorem 6.4, and it is equal to $A_\alpha^p(\mathbb{B}^n)$. Since u is smooth vector for $\mathcal{H}_s$, and $\mathcal{H}_s^\infty$ is continuously embedded in its dual $\mathcal{H}_s^{-\infty}$, the vector u is ρ - and ρ^* -weakly differentiable. According to Theorem 6.6, the mappings

$$f \mapsto f * |W_{\rho(E_\alpha)u}^{\rho_s}(u)| \quad \text{and} \quad f \mapsto f * |W_u^{\rho_s}(\rho^*(E_\alpha)u)|$$

are continuous on $L_{\alpha+n+1-ps/2}^p(G)$. Therefore, we can choose ϵ small enough such that the family $\{\rho_s(x_i)u\}$ forms a frame and an atomic decomposition for $A_\alpha^p(\mathbb{B}^n)$ with reconstruction operators that are given in Theorem 5.3. $\square$

APPENDIX A. DECOMPOSITIONS OF REPRODUCING KERNEL SPACES FOR TWISTED CONVOLUTION

From now on we let $\phi(x) = W_u^\rho(u)$ for some fixed $u \in \mathcal{S}$. Then $B_u^\sigma = \{f \in B \mid f = f \# \phi\}$. Given a compact neighbourhood U of the identity in G , a U -dense and well-spread sequence $\{x_i\} \subseteq G$ and an U -BUPU $\{\psi_i\}$, we formally define the

operators

$$\begin{aligned} T_1 f &= \sum_i f(x_i) \sigma(x, x^{-1} x_i) \psi_i \# \phi \\ T_2 f &= \sum_i \lambda_i(f) \ell_{x_i}^\sigma \phi \\ T_3 f &= \sum_i c_i f(x_i) \ell_{x_i}^\sigma \phi \end{aligned}$$

where $\lambda_i(f) = \int f(y) \psi_i(y) \overline{\sigma(y, y^{-1} x_i)} dy$ and $c_i = \int \psi_i(y) dy$. The following results will establish when these operators are well defined on B_u^σ .

Define the local oscillations

$$\text{osc}_U^r f(x) = \sup_{y \in U} |r_y^\sigma f(x) - f(x)| \quad \text{and} \quad \text{osc}_U^{\ell^\sigma} f(x) = \sup_{y \in U} |\ell_y^\sigma f(x) - f(x)|.$$

Proposition A.1. *If $f \in B_u^\sigma$ then*

$$\begin{aligned} |T_1 f(x) - f(x)| &\leq |f| * \text{osc}_{U^{-1}}^r \phi(x) \\ |T_2 f(x) - f(x)| &\leq |f| * \text{osc}_{U^{-1}}^{\ell^\sigma} \phi(x) \\ |T_3 f(x) - f(x)| &\leq |f| * \text{osc}_{U^{-1}}^r \phi * (|\phi| + \text{osc}_{U^{-1}}^{\ell^\sigma} \phi)(x) + |f| * \text{osc}_{U^{-1}}^{\ell^\sigma} \phi(x). \end{aligned}$$

Proof. We see that

$$\left| \sum_i f(x_i) \sigma(x, x^{-1} x_i) \psi_i(x) - f(x) \right| \leq \sum_i |f(x_i) \sigma(x, x^{-1} x_i) - f(x)| \psi_i(x)$$

and for $x \in x_i U$ we get that $x_i \in x U^{-1}$, so

$$|f(x_i) \sigma(x, x^{-1} x_i) - f(x)| \leq \sup_{y \in U^{-1}} |f(xy) \sigma(x, y) - f(x)| = \text{osc}_{U^{-1}}^r f(x).$$

Next, if $f = f \# \phi$ we get

$$|r_y^\sigma f(x) - f(x)| \leq \int |f(z)| |\phi(z^{-1} xy) \overline{\sigma(z, z^{-1} xy)} \sigma(x, y) - \phi(z^{-1} x) \overline{\sigma(z, z^{-1} x)}| dz.$$

Since $\sigma(z, z^{-1} xy) = \sigma(z, z^{-1} x) \sigma(x, y) \overline{\sigma(z^{-1} x, y)}$ we get that the integral above reduces to

$$\int |f(z)| |r_y^\sigma \phi(z^{-1} x) - \phi(z^{-1} x)| dz.$$

Taking supremum we get the desired result.

We have that

$$\begin{aligned} f(x) - T_2 f(x) &= \int f(y) \ell_y^\sigma \phi(x) dy - \int \sum_i f(y) \psi_i(y) \overline{\sigma(y, y^{-1} x_i)} dy \ell_{x_i}^\sigma \phi(x) \\ &= \int \sum_i f(y) \psi_i(y) [\ell_y^\sigma \phi(x) - \overline{\sigma(y, y^{-1} x_i)} \ell_{x_i}^\sigma \phi(x)] dy \\ &= \int \sum_i f(y) \psi_i(y) \ell_y^\sigma [\phi(x) - \ell_{y^{-1} x_i}^\sigma \phi(x)] dy. \end{aligned}$$

When $y \in x_i U$ for a compact neighbourhood U of the identity, then $y^{-1}x_i \in U^{-1}$. Thus we get

$$\begin{aligned} |f(x) - T_2 f(x)| &\leq \int \sum_i |f(y)| |\psi_i(y)| |\ell_y^\sigma \phi(x) - \ell_{y^{-1}x_i}^\sigma \phi(x)| dy \\ &\leq \int \sum_i |f(y)| |\psi_i(y)| |\ell_y^\sigma \text{osc}_{U^{-1}}^{\ell^\sigma} \phi| dy \\ &\leq \int |f(y)| \ell_y \text{osc}_{U^{-1}}^{\ell^\sigma} \phi dy. \end{aligned}$$

This shows the claim.

$$|T_3 f(x) - f(x)| \leq \int \sum_i \psi_i(y) |f(x_i) \ell_{x_i}^\sigma \phi(x) - f(y) \ell_y^\sigma \phi(x)| dy$$

Let us rewrite part of the integrand when $y \in x_i U$

$$\begin{aligned} &|f(x_i) \ell_{x_i}^\sigma \phi(x) - f(y) \ell_y^\sigma \phi(x)| \\ &= |f(x_i) \sigma(y, y^{-1}x_i) \ell_y^\sigma \ell_{y^{-1}x_i}^\sigma \phi(x) - f(y) \ell_y^\sigma \phi(x)| \\ &\leq |[f(x_i) \sigma(y, y^{-1}x_i) - f(y)] \ell_y^\sigma \ell_{y^{-1}x_i}^\sigma \phi(x)| + |f(y)| |\ell_y^\sigma \ell_{y^{-1}x_i}^\sigma \phi(x) - \ell_y^\sigma \phi(x)| \\ &= |[r_{y^{-1}x_i}^\sigma f(y) - f(y)] \ell_y^\sigma \ell_{y^{-1}x_i}^\sigma \phi(x)| + |f(y)| |\ell_y^\sigma \ell_{y^{-1}x_i}^\sigma \phi(x) - \ell_y^\sigma \phi(x)| \\ &\leq \text{osc}_{U^{-1}}^\sigma f(y) [|\ell_y^\sigma \ell_{y^{-1}x_i}^\sigma \phi(x) - \ell_y^\sigma \phi(x)| + |\ell_y^\sigma \phi(x)|] + |f(y)| |\ell_y^\sigma \text{osc}_{U^{-1}}^{\ell^\sigma} \phi(x)| \\ &\leq \text{osc}_{U^{-1}}^\sigma f(y) [|\ell_y^\sigma \text{osc}_{U^{-1}}^{\ell^\sigma} \phi(x)| + |\ell_y^\sigma \phi(x)|] + |f(y)| |\ell_y^\sigma \text{osc}_{U^{-1}}^{\ell^\sigma} \phi(x)|. \end{aligned}$$

As before the oscillation of f can be transferred onto the kernel, and the final result is obtained. $\square$

From this we obtain

Corollary A.2. *Let B be a solid BF-space and assume that $f \mapsto f * |\phi|$, $f \mapsto f * \text{osc}_{U^{-1}}^{\ell^\sigma} \phi$ and $f \mapsto f * \text{osc}_{U^{-1}}^{\ell^\sigma} \phi$ are bounded on B , then T_1, T_2, T_3 are well-defined bounded operators on $B_u^\#$.*

Moreover, if there are constants C_U for which

$$\|f * \text{osc}_{U^{-1}}^{\ell^\sigma} \phi\| \leq C_U \|f\| \text{ and } \|f * \text{osc}_{U^{-1}}^{\ell^\sigma} \phi\| \leq C_U \|f\|$$

and $\lim_{U \rightarrow \{e\}} C_U = 0$, then there is a U small enough as well as U -dense $\{x_i\}$ such that the operators are invertible on $B_u^\#$.

We will now use the special form of ϕ to find oscillation estimates via derivatives. We have defined $\phi(x) = \langle u, \rho(x)u \rangle$, and from this and Remark 3.2 we get

$$\text{osc}_U^\sigma \phi(x) = \sup_{y \in U} |\langle \rho^*(x^{-1})u, \rho(y)u - u \rangle|,$$

and

$$\text{osc}_U^{\ell^\sigma} \phi(x) = \sup_{y \in U} |\langle \rho^*(y)u - u, \rho(x)u \rangle|.$$

In light of this it seems possible to evaluate the oscillation by a certain level of smoothness of the vector u , and this is exactly the approach we will take. We let $v \in \mathcal{S}$ and $\lambda \in \mathcal{S}^*$ be arbitrary elements and define the functions $H(y) = \langle \lambda, \rho(y)u \rangle$ and $K(y) = \langle \rho^*(y)u, v \rangle$. We will now investigate the local oscillations of H and K in terms of derivatives, but first we need to introduce some notation.

If f is a function on G and X is in $\mathfrak{g}$, then define

$$Xf(y) = \frac{d}{dt} \Big|_{t=0} f(y \exp(tX)).$$

We now fix a basis $X_1, \dots, X_n$ for the Lie algebra $\mathfrak{g}$, and for a multi-index α we define

$$X^\alpha f = X_1^{\alpha_1} \dots X_n^{\alpha_n} f.$$

We will investigate oscillations of H and K on the specific neighbourhood

$$U_\epsilon = \{\exp(t_1 X_1) \dots \exp(t_n X_n) \mid -\epsilon \leq t_k \leq \epsilon\}.$$

Remember that we choose the cocycle σ and $\epsilon > 0$ such that σ is C^∞ on a neighbourhood containing $U_\epsilon \times U_\epsilon$. According to Lemma 2.5 in [5] there is a constant C_ϵ such that

$$\sup_{y \in U_\epsilon} |H(y) - H(e)| \leq C_\epsilon \sum_{\substack{1 \leq |\alpha| \leq n \\ |\delta| = |\alpha|}} \int_{[-\epsilon, \epsilon]^{|\delta|}} |X^\alpha H(\tau_\delta(t_1, \dots, t_n))| (dt_1)^{\delta_1} \dots (dt_n)^{\delta_n},$$

and

$$\sup_{y \in U_\epsilon} |K(y) - K(e)| \leq C_\epsilon \sum_{\substack{1 \leq |\alpha| \leq n \\ |\delta| = |\alpha|}} \int_{[-\epsilon, \epsilon]^{|\delta|}} |X^\alpha K(\tau_\delta(t_1, \dots, t_n))| (dt_1)^{\delta_1} \dots (dt_n)^{\delta_n},$$

where $\tau_\delta(t_1, \dots, t_n) = \exp(\delta_1 t_1 X_1) \dots \exp(\delta_n t_n X_n)$ for a multi-index δ . Due to the special form of H

$$XH(y) = \frac{d}{dt} \Big|_{t=0} \langle \lambda, \rho(y \exp(tX))u \rangle = \frac{d}{dt} \Big|_{t=0} \langle \lambda, \rho(y) \rho(\exp(tX))u \rangle \overline{\sigma(y, \exp(tX))}.$$

Therefore $X^\delta H(y)$ can be expressed as a sum

$$\sum_{|\gamma| \leq |\delta|} \langle \lambda, \rho(y) \rho(X_n)^{\gamma_n} \dots \rho(X_1)^{\gamma_1} u \rangle g_\gamma(y),$$

where g_γ is an appropriate derivative of the cocycle σ of order $|\gamma|$. Notice, that the g_γ 's do not depend on the vectors v and λ used to define H and K . If y is in the compact set U_ϵ the functions g_γ are uniformly bounded, and therefore there is a constant D_ϵ such that

$$\begin{aligned} & \sup_{y \in U_\epsilon} |H(y) - H(e)| \\ & \leq D_\epsilon \sum_{\substack{1 \leq |\alpha| \leq n \\ |\delta| = |\alpha|}} \int_{[-\epsilon, \epsilon]^{|\delta|}} |\langle \lambda, \rho(\tau_\delta(\mathbf{t})) \rho(X_n)^{\delta_n} \dots \rho(X_1)^{\delta_1} u \rangle| (dt_1)^{\delta_1} \dots (dt_n)^{\delta_n}, \end{aligned}$$

when we write $\mathbf{t} = (t_1, t_2, \dots, t_n)$. From this we get that

$$\begin{aligned} & \text{osc}_U^{\sigma} \phi(x) \\ & \leq D_\epsilon \sum_{\substack{1 \leq |\alpha| \leq n \\ |\delta| = |\alpha|}} \int_{[-\epsilon, \epsilon]^{|\delta|}} |\langle u, \rho(x \tau_\delta(\mathbf{t})) \rho(X_n)^{\delta_n} \dots \rho(X_1)^{\delta_1} u \rangle| (dt_1)^{\delta_1} \dots (dt_n)^{\delta_n}. \end{aligned}$$

Treating K the same way we get

$$\begin{aligned} & \text{osc}_U^{\ell^\sigma} \phi(x) \\ & \leq D_\epsilon \sum_{\substack{1 \leq |\alpha| \leq n \\ |\delta| = |\alpha|}} \int_{[-\epsilon, \epsilon]^{|\delta|}} |\langle \rho^*(X_n)^{\delta_n} \cdots \rho^*(X_1)^{\delta_1} u, \rho(\tau_\delta(\mathbf{t})^{-1}x)u \rangle| (dt_1)^{\delta_1} \cdots (dt_n)^{\delta_n}. \end{aligned}$$

Lemma A.3. *Assume that B is a solid BF-space on which left and right translations are continuous. If $f \mapsto f * |\langle u, \rho(\cdot)\rho(X_n)^{\delta_n} \cdots \rho(X_1)^{\delta_1} u \rangle|$ and $f \mapsto f * |\langle \rho^*(X_n)^{\delta_n} \cdots \rho^*(X_1)^{\delta_1} u, \rho(\cdot)u \rangle|$ are bounded on B for all $|\delta| \leq \dim(G)$, then*

$$\|f * \text{osc}_U^{\ell^\sigma} \phi\|_B \leq C_\epsilon \|f\|_B$$

and

$$\|f * \text{osc}_U^{r^\sigma} \phi\|_B \leq C_\epsilon \|f\|_B.$$

Moreover, $\lim_{\epsilon \rightarrow 0} C_\epsilon = 0$.

Proof. Write $\mathbf{t} = (t_1, \dots, t_n)$. Notice that

$$\begin{aligned} & |f * \text{osc}_U^{\ell^\sigma} \phi(x)| \\ & \leq D_\epsilon \sum_{\substack{1 \leq |\alpha| \leq n \\ |\delta| = |\alpha|}} \int_{[-\epsilon, \epsilon]^{|\delta|}} |\ell_{\tau_\delta(\mathbf{t})} f| * |W_{\rho^*(X_n)^{\delta_n} \cdots \rho^*(X_1)^{\delta_1} u}(u)(x)| (dt_1)^{\delta_1} \cdots (dt_n)^{\delta_n}. \end{aligned}$$

Since left translation is continuous on B the right hand side defines a function in B by Theorem 3.29 in [30], and by solidity $f * \text{osc}_U^{\ell^\sigma} \phi(x)$ is also in B . Moreover, $\|f * \text{osc}_U^{\ell^\sigma} \phi\| \leq C_\epsilon \|f\|$, where C_ϵ is equal to D_ϵ multiplied by a polynomial in ϵ with no constant term. Since D_ϵ is uniform in ϵ we see that $\lim_{\epsilon \rightarrow 0} C_\epsilon = 0$.

The proof for convolution with right oscillations follows in a similar manner. $\square$

REFERENCES

- [1] P. Aniello. Square integrable projective representations and square integrable representations modulo a relatively central subgroup. *Int. J. Geom. Methods Mod. Phys.*, 3(2):233–267, 2006.
- [2] V. Bargmann. On unitary ray representations of continuous groups. *Ann. of Math. (2)*, 59:1–46, 1954.
- [3] H. Chébli and J. Faraut. Fonctions holomorphes à croissance modérée et vecteurs distributions. *Math. Z.*, 248(3):540–565, 2004.
- [4] J. Christensen, K. Gröchenig, and G. Ólafsson. New atomic decompositions for Bergman spaces on the unit ball. *ArXiv e-prints*, Apr. 2015.
- [5] J. G. Christensen. Sampling in reproducing kernel Banach spaces on Lie groups. *Journal of Approximation Theory*, 164(1):179–203, 2012.
- [6] J. G. Christensen, A. Mayeli, and G. Ólafsson. Coorbit Description and Atomic Decomposition of Besov Spaces. *Numerical Functional Analysis and Optimization*, 33(7-9):847–871, 2012.
- [7] J. G. Christensen and G. Ólafsson. Coorbit spaces for dual pairs. *Appl. Comput. Harmon. Anal.*, 31(2):303–324, 2011.
- [8] J. G. Christensen and G. Ólafsson. Sampling in spaces of bandlimited functions on commutative spaces. In *Excursions in harmonic analysis. Volume 1*, Appl. Numer. Harmon. Anal., pages 35–69. Birkhäuser/Springer, New York, 2013.
- [9] O. Christensen. Atomic decomposition via projective group representations. *Rocky Mountain J. Math.*, 26(4):1289–1312, 1996.
- [10] R. R. Coifman and R. Rochberg. Representation theorems for holomorphic and harmonic functions in L^p . In *Representation theorems for Hardy spaces*, volume 77 of *Astérisque*, pages 11–66. Soc. Math. France, Paris, 1980.

- [11] S. Dahlke, F. De Mari, E. De Vito, D. Labate, G. Steidl, G. Teschke, and S. Vigogna. Coorbit Spaces with Voice in a Fréchet Space. *Journal of Fourier Analysis and Applications*, pages 1–66, 2016.
- [12] S. Dahlke, M. Fornasier, H. Rauhut, G. Steidl, and G. Teschke. Generalized coorbit theory, Banach frames, and the relation to α -modulation spaces. *Proc. Lond. Math. Soc. (3)*, 96(2):464–506, 2008.
- [13] S. Dahlke, G. Steidl, and G. Teschke. Coorbit spaces and Banach frames on homogeneous spaces with applications to the sphere. *Adv. Comput. Math.*, 21(1-2):147–180, 2004.
- [14] S. Dahlke, G. Steidl, and G. Teschke. Weighted coorbit spaces and Banach frames on homogeneous spaces. *J. Fourier Anal. Appl.*, 10(5):507–539, 2004.
- [15] S. Dahlke, G. Steidl, and G. Teschke. Frames and coorbit theory on homogeneous spaces with a special guidance on the sphere. *J. Fourier Anal. Appl.*, 13(4):387–404, 2007.
- [16] A. H. Darweesh. *Wavelets, Coorbit Theory, and Projective Representation*. PhD thesis, Louisiana State University, LSU, 2015.
- [17] J. Faraut and A. Korányi. Function spaces and reproducing kernels on bounded symmetric domains. *J. Funct. Anal.*, 88(1):64–89, 1990.
- [18] H. G. Feichtinger and K. Gröchenig. A unified approach to atomic decompositions via integrable group representations. In *Function spaces and applications (Lund, 1986)*, volume 1302 of *Lecture Notes in Math.*, pages 52–73. Springer, Berlin, 1988.
- [19] H. G. Feichtinger and K. Gröchenig. Banach spaces related to integrable group representations and their atomic decompositions. I. *J. Funct. Anal.*, 86(2):307–340, 1989.
- [20] H. G. Feichtinger and K. Gröchenig. Banach spaces related to integrable group representations and their atomic decompositions. II. *Monatsh. Math.*, 108(2-3):129–148, 1989.
- [21] M. Fornasier and H. Rauhut. Continuous frames, function spaces, and the discretization problem. *J. Fourier Anal. Appl.*, 11(3):245–287, 2005.
- [22] K. Gröchenig. Describing functions: atomic decompositions versus frames. *Monatsh. Math.*, 112(1):1–42, 1991.
- [23] Harish-Chandra. Representations of semisimple Lie groups. IV. *Amer. J. Math.*, 77:743–777, 1955.
- [24] A. W. Knap. *Representation theory of semisimple groups*, volume 36 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1986. An overview based on examples.
- [25] D. H. Luecking. Representation and duality in weighted spaces of analytic functions. *Indiana Univ. Math. J.*, 34(2):319–336, 1985.
- [26] G. W. Mackey. Unitary representations of group extensions. I. *Acta Math.*, 99:265–311, 1958.
- [27] N. S. Poulsen. On *C*-spinfty*-vectors and intertwining bilinear forms for representations of Lie groups. *J. Functional Analysis*, 9:87–120, 1972.
- [28] H. Rauhut. *Time-Frequency and Wavelet Analysis of Functions with Symmetry Properties*. Logos-Verlag, 2005.
- [29] H. Rauhut. Coorbit space theory for quasi-Banach spaces. *Studia Math.*, 180(3):237–253, 2007.
- [30] W. Rudin. *Functional analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill Inc., New York, second edition, 1991.
- [31] V. S. Varadarajan. *Geometry of quantum theory*. Springer-Verlag, New York, second edition, 1985.
- [32] N. R. Wallach. The analytic continuation of the discrete series. I, II. *Trans. Amer. Math. Soc.*, 251:1–17, 19–37, 1979.
- [33] G. Warner. *Harmonic analysis on semi-simple Lie groups. I*. Springer-Verlag, New York, 1972.
- [34] K. Zhu. *Spaces of holomorphic functions in the unit ball*, volume 226 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.

DEPARTMENT OF MATHEMATICS, COLGATE UNIVERSITY, 13 OAK DRIVE, HAMILTON NY 13346

E-mail address: `jchristensen@colgate.edu`

URL: `http://www.math.colgate.edu/~jchristensen`

DEPARTMENT OF MATHEMATICS AND STATISTICS, JORDAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, IRBID 22110, B.O. BOX (3030) JORDAN.

E-mail address: `ahdarweesh@just.edu.jo`

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803

E-mail address: `olafsson@math.lsu.edu`

URL: `http://www.math.lsu.edu/~olafsson`